




SPRING AESTHETICS SALE









To Order: 1-800-P-SCHEIN (1-800-772-4346) 8am-9pm (et)

Spring Aesthetics Sale

EXPAREL®

(bupivacaine liposome injectable suspension) for Postsurgical Pain Control



For important safety information and to learn more about Exparel, go to:

www.exparel.com/index.shtml

For medical inquiries related to EXPAREL, contact Medical Information at medinfo@pacira.com or 1-855-RX-EXPAREL (1-855-793-9727).

*Pivotal studies have demonstrated the safety and efficacy of EXPAREL in patients undergoing bunionectomy or hemorrhoidectomy procedures; additional studies are underway to further demonstrate the safety and efficacy in other procedures.

†The clinical benefit of the attendant decrease in opioid consumption was not demonstrated.

¹EXPAREL [prescribing information]. Parsippany, NJ: Pacira Pharmaceuticals, Inc.; 2011.

²Gorfine SR, Onel E, et al. Bupivacaine extended-release liposome injection for prolonged postsurgical analgesia in patients undergoing hemorrhoidectomy: a multicenter, randomized, double-blind, placebo-controlled trial. Diseases of the Colon & Rectum. 2011; 54(12):1559.

What is Exparel?

EXPAREL®—Postsurgical Pain Control That Starts When Surgery Ends and Continues Days Into Recoverv*

EXPAREL® is a local analgesic that utilizes bupivacaine in combination with the proven product delivery platform, DepoFoam®. A single intraoperative injection given at the close of surgery delivers postsurgical pain control with reduced opioid requirements† for up to 72 hours.**1

EXPAREL® is the first and only multivesicular liposome-based local anesthetic that can be utilized in the peri- or postsurgical setting in the same fashion as current local anesthetics. Local infiltration of EXPAREL® results in significant systemic plasma levels of bupivacaine that can persist for 96 hours.

- Systemic plasma levels of bupivacaine following administration of EXPAREL® are not correlated with local efficacy
- Other formulations of bupivacaine should not be administered within 96 hours following administration of EXPAREL®

EXPAREL® is indicated for administration into the surgical site to produce postsurgical analgesia.¹

The EXPAREL® Difference

Same Infiltration Technique Provides Several Days of Postsurgical Analgesia.* Surgeons who infiltrate local anesthetics can use the same technique to administer EXPAREL® at the close of surgery. As a non-opioid, single-dose analgesic¹, EXPAREL® can:

- Provide up to 72 hours* of analgesia with reduced opioid requirements[†]
- Without the need for catheters or pumps

In a pivotal hemorrhoidectomy trial of EXPAREL® compared to placebo, where all patients with inadequate pain control received opioids for rescue pain relief, a single intraoperative injection of EXPAREL® demonstrated:

- A 30% reduction in cumulative pain scores and more pain-free patients at all time points up to and including 24 hours (P<0.0001)²
- A 45% decrease in the use of opioid rescue medication at 72 hours and more than a 12-fold increase in the median time to first opioid use (P=0.0006 and P<0.0001, respectively)²; the clinical benefit of the attendant decrease in opioid consumption was not determined

EXPAREL® is contraindicated in obstetrical paracervical block anesthesia. EXPAREL® has not been studied for use in patients younger than 18 years of age.

Spring Aesthetics Sale



Bacitracin Powder for Injection

Compare to Baciguent® 50,000 U, 30 mL (908-8136)ea



Botox® Cosmetic Injectable 100-U Vial

(377-6699)ea



Celestone® Soluspan® Injectable Suspension 6 mg/mL, 5-mL MDV

(516-0266)ea



Compare & SAVE!

Betamethasone Sodium Phosphate and **Betamethasone Acetate**

Celestone® Soluspan® 6 mg/mL, 5-mL MDV (114-8668)ea



Drysol Solution, 20% 37.5-mL/Bottle

(928-6031)ea





Great

Hylenex (Human)

Hylenex® recombinant (hyaluronidase human injection). 1-mL Injectable, 150 U/0mL SDV (248-0470) 4/pkg



Kenalog®-10 Injectable Triamcinolone Acetonide 10 mg/mL, 5-mL MDV

(196-0545)ea



Kenalog®-40 Injectables

Triamcinolone Acetonide 40 mg/mL, 5-mL MDV

(196-1737)ea 40 mg/mL, 10-mL MDV (196-9429)ea





Lidocaine HCI Injectable

Compare to Xylocaine® 50-mL MDVs 1%

(104-6817)25/box





Lidocaine with Epinephrine 1:100 Injectable

Compare to Xylocaine® with Epinephrine 30-mL MDV 1% (104-6822)25/box





Lidocaine 1% with Epinephrine 1:100 Injectable

Compare to Xylocaine®

with Epinephrine 50-mL MDV (104-7099) 25/box



Mupirocin 2% Ointment

Compare to Bactroban® 22-g Tube (111-8857).....ea



Sevoflurane Liquid Inhalation 100% 250 mL/Bottle

(116-9356)ea



Sotradecol Injectable

1%, 10 mg/mL, 2-mL Vial (424-0048)......5/pkg 3%, 30 mg/mL, 2-mL Vial **(424-0049)**......5/pkg



Xylocaine® with Epinephrine Injectable

1%, 50-mL MDV (248-0350).....ea 2%, 50-mL MDV (248-0414)ea

To Fax an Order

Mention promo code SZ to ensure promotional pricing Offer Ends June 30, 2013

HENRY SCHEIN, INC.

J.S. POSTAGE PRSRT STD

Henry Schein, Inc. • 135 Duryea Road • Melville, NY 11747



Exclusive!

A versatile, professional chemical peel hygienically packaged in a full procedure kit



Average Physician Fee \$150±\$250

For complete product information or to order, visit www.henryschein.com/bliftx

Complete Kit:

B-Liftx® is packaged for a successful patient experience and contains the following:

- 2 alcohol precleanse pads
- 1 vial of either 20% or 30% salicylic acid in an alcohol solution
- Dispensing cup and applicator sponge
- 1 tube of 0.75 oz. HydroLotion® for post-procedure skin care
 - Uniquely formulated with special petrolatum fraction to provide deep, long-lasting moisturization
 - Noncomedogenic, hypoallergenic, fragrance and paraben-free

Each kit is hygienically packed for single patient use and packed 24 per case

20% Salicylic Acid Peel Kits (118-7633)

30% Salicylic Acid Peel Kits (118-7632)

β-Liftx® is a registered trademark of DermThink LLC and is used under license. All rights reserved. No mix and match. Free goods ship with order.