

Dermabond Advanced®

DERMABOND ADVANCED®
Topical Skin Adhesive
Evidence Brief



Overview

As the final layer of wound closure, topical skin adhesives (TSAs) are an integral part of a successful clinical outcome. When deciding which TSA to use, clinical study information on closure strength, microbial protection, patient comfort, and cosmesis allows healthcare practitioners to evaluate which product will provide the greatest benefits for their patients.

DERMABOND ADVANCED® Topical Skin Adhesive is supported by an extensive body of published literature, including 53 randomized controlled trials (RCTs). DERMABOND ADVANCED Adhesive has a patented, proprietary chemical formulation¹ that has been shown to provide superior strength versus other commercially available TSAs,² and also has benefits that enhance patient comfort and cosmetic outcomes.³⁻⁶

This Evidence Summary includes a sample of the available RCTs for DERMABOND ADVANCED Adhesive or DERMABOND® Topical Skin Adhesive. **A full list of published studies can be found in the bibliography section of this document.**

- DERMABOND ADVANCED Adhesive and DERMABOND Adhesive are supported by 53 published RCTs*†
- Total of 5,836 patients evaluated

References

1. DERMABOND ADVANCED® Topical Skin Adhesive Label. LAB-0012182 DNX12. Ethicon, Inc.
2. Singer AJ, Perry LC, Allen RL. In vivo study of wound-bursting strength and compliance of topical skin adhesives. *Acad Emerg Med*. 2008;15(12):1290-1294.
3. Nipshagen MD, Hage JJ, Beekman W. Use of 2-octyl-cyanoacrylate skin adhesive (Dermabond) for wound closure following reduction mammoplasty: a prospective, randomized intervention study. *Plast Reconstr Surg*. 2008;122:10-18.
4. Scott GR, Carson CL, Borah G. Dermabond skin closures for bilateral reduction mammoplasties: a review of 255 consecutive cases. *Plast Reconstr Surg*. 2007;120:1460-1465.
5. Toriumi DM, O'Grady K, Desai D, Bagal A. Use of octyl-2-cyanoacrylate for skin closure in facial plastic surgery. *Plast Reconstr Surg*. 1998;102:2209-2219.
6. Quinn J, Wells G, Sutcliffe T, et al. A randomized trial comparing octylcyanoacrylate tissue adhesive and sutures in the management of lacerations. *JAMA*. 1997;277(19):1527-1530.

*DERMABOND ADVANCED Adhesive tests equivalent or superior to DERMABOND Adhesive in head-to-head testing for microbial barrier, wound-bursting strength, tensile strength, flexibility, durability, viscosity, drying time, water vapor transmission rate, water resistance, and physician satisfaction.

†Based on published literature in PubMed and SCOPUS, using only RCTs that evaluated the use of the product in a manner consistent with intended indication.

Summary of Key Studies

The publications that support the claims for DERMABOND ADVANCED® Topical Skin Adhesive are listed in the table below. A summary of each of these studies can be found on the subsequent pages.

Publication Title	Lead Author	Source	Outcome Studied
In vitro Assessment of Microbial Barrier Properties of DERMABOND® Topical Skin Adhesive	Bhende	<i>Surgical Infections.</i> 2002;3(3):251-257.	Microbial Barrier
In vitro study to determine the ability of DERMABOND ADVANCED® Topical Skin Adhesive to inhibit bacterial growth	Bhende	Internal Ethicon Study	Inhibition of Bacteria
In Vivo Study of Wound Bursting Strength and Compliance of Topical Skin Adhesives	Singer	<i>Academic Emergency Medicine.</i> 2008;15(12):1290-1294.	Strength and Flexibility
Postoperative Outcomes Associated with Topical Skin Adhesives among Women Having Hysterectomies	Murmann	<i>Surgical Infections.</i> 2010;11(5):441-447.	Hospitalization Costs
A Randomized Trial Comparing Octylcyanoacrylate Tissue Adhesive and Sutures in the Management of Lacerations	Quinn	<i>JAMA.</i> 1997;277(19):1527-1530.	Cosmesis, Time, Pain

Clinical Reference Article Summary

In Vitro Assessment of Microbial Barrier Properties of DERMABOND® Topical Skin Adhesive

Bhende S, Rothenburger S, Spangler D, Dito M

Source:

Surgical Infections. 2002;3(3):251-257

Study Objective

The purpose of this study was to evaluate the ability of DERMABOND Adhesive to provide an effective microbial barrier against the penetration of microorganisms in vitro.

Bacteria used in this study included:

Staphylococcus aureus

Staphylococcus epidermidis

Escherichia coli

Pseudomonas aeruginosa

Enterococcus faecium

Methods

Plates containing an agar media were created in a sterile environment. The agar media contained a pH-sensitive dye designed to color when exposed to the acidic metabolic products of bacteria.

DERMABOND Adhesive was applied to the agar surface. In total, 300 single-layer films and 300 triple-layer films were examined. The surface of each film was inoculated with a 10 µL aliquot of bacteria containing at least 1×10^3 colony-forming units (cfu).

All test and control plates were incubated at 37°C for 72 hours. A change in color indicated a breach in the adhesive's microbial barrier.

Results

Single-layer films: 299 of the 300 samples retained their integrity as microbial barriers for 72 hours.

All 300 samples maintained their microbial barrier for 48 hours.

For the triple-layer films, 299 of the 300 samples retained their integrity as microbial barriers for 72 hours.

Conclusion

The results of this study demonstrate that DERMABOND Adhesive provides a microbial barrier with 99% protection in vitro for at least 72 hours against organisms commonly responsible for SSIs, including: *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Enterococcus faecium*.

Clinical Reference Article Summary

In vitro study to determine the ability of DERMABOND ADVANCED® Topical Skin Adhesive to inhibit bacterial growth

Bhende S

Source:

Internal Ethicon Study

Study Objective

The purpose of this study was to demonstrate that DERMABOND ADVANCED Adhesive inhibits gram-positive bacteria and gram-negative bacteria in vitro.

Bacteria evaluated in this study:

Methicillin-resistant *Staphylococcus aureus* (MRSA)

Methicillin-resistant *Staphylococcus epidermidis* (MRSE)

Escherichia coli

Methods

Cultures of each organism were grown under sterile conditions for 18-24 hours at 35-37°C. Before being used in the experiment, each culture was diluted to achieve an approximate bacteria count of 10⁵ colony-forming units (cfu)/0.04 ml.

A 2 cm diameter circle was drawn on the bottom of a sterile agar plate. In the center of this circle, 0.04 ml of the diluted inoculum was placed on the surface of the agar.

After allowing the inoculum to dry, the adhesive material was applied to the inoculated surface area, making sure to cover the area beyond the marked circle.

After 10 minutes of contact time between the adhesive and the inoculated area, the adhesive's polymerized film was removed from the surface of the agar, and the plates were incubated at 37°C for up to 48 hours.

In total, 210 samples (70 samples per organism) were evaluated. The samples were examined for bacterial growth at 24 and 48 hours. Any growth originating beneath the area of adhesive application was recorded as a positive test.

Results

After 48 hours, the test plates exhibited colony counts ranging from 0 - 59 cfu, indicating significant inhibition of the bacteria.

Each inoculated plate was declared a success if a minimum of 99.9% inhibition of the initial inoculum load was observed. For all bacteria evaluated (MRSA, MRSE, *E. coli*), contact with the adhesive led to a 99.9% inhibition in bacteria load from the initial inoculum.

Conclusion

In this in vitro study, DERMABOND ADVANCED Adhesive was shown to demonstrate inhibition of gram-positive bacteria (MRSA, MRSE) and gram-negative bacteria (*E. coli*)*.

*Clinical significance is unknown.

Clinical Reference Article Summary

In Vivo Study of Wound Bursting Strength and Compliance of Topical Skin Adhesives

Singer AJ, Perry LC, Allen Jr. RL

Source:

Academic Emergency Medicine. 2008;15(12):1290-1294

Study Objective

The purpose of this study was to evaluate the wound-bursting strength and flexibility of five topical skin adhesives during the two-day period after wound closure.

The following adhesives were evaluated in the study:

DERMABOND® Topical Skin Adhesive

INDERMIL® Tissue Adhesive

Histoacryl® Topical Skin Adhesive

LiquiBand® Topical Skin Adhesive

GluStitch®

Methods

Using a template for incision length and location, two symmetric incisions (2 cm long each) were created over the dorsolateral flank area of 210 anesthetized, male Sprague-Dawley rats.

After achieving hemostasis and manually approximating the skin edges, a randomized computer algorithm was used to select an adhesive to close the incision. All adhesives were applied according to manufacturer instructions.

The adhesives were evaluated three times during the study—immediately after closure, 1 day after closure, and 2 days after closure.


For each evaluation, 14 samples from each adhesive group were tested for wound-bursting strength, and another 14 samples were tested for flexibility.

To test for wound-bursting strength, a vacuum chamber was placed over each sample and negative pressure was applied, stressing the wound in 3 dimensions. The pressure (mmHg) needed to cause wound failure was recorded.

To test for flexibility, a vacuum chamber was placed over the sample and negative pressure was applied to the wound while a laser measured the vertical deformation of the skin (μm). Energy absorption (mmHg x mm) was calculated to quantify the adhesives' flexibility.


Results

Figure 1
Wound-bursting Strength



In total, 210 measurements were taken on 210 incisions (5 adhesives, 3 time points, 14 samples per time point). Results are shown in **Figure 1**.


Figure 2
Percent of Samples with Visible Fractures



With the exception of the samples in the DERMABOND® Topical Skin Adhesive group, measurements could not be taken on all samples in an adhesive group because, in some samples, the adhesive's inflexibility had caused the adhesive to fracture during testing.

As shown in **Figure 2**, the percent of samples in an adhesive group experiencing fractures ranged from 36% to 86%.

Figure 3
Flexibility of Five Topical Skin Adhesives



As seen in **Figure 3**, for the samples that maintained their integrity throughout testing, the samples in the DERMABOND Adhesive group consistently had the greatest flexibility. Adhesive flexibility decreased over time in all cases.

Conclusion

The results of this study demonstrate that DERMABOND Adhesive was significantly stronger and more flexible than the other adhesives evaluated in the study.*

*This study was funded in full or in part by an educational grant from Ethicon, Inc.

Clinical Reference Article Summary

Postoperative Outcomes Associated with Topical Skin Adhesives among Women Having Hysterectomies

Murrmann SG, Markowitz JS, Gutterman EM, Magee G

Source:

Surgical Infections. 2010;11(5):441-447

Study Objective

The purpose of this study was to evaluate the clinical and economic outcomes associated with use of a topical skin adhesive (TSA) versus traditional methods for skin closure following total abdominal hysterectomy.

Methods

The study utilized Premier Perspective™ Comparative Database, which is a large, administrative database containing clinical and economic data from all patient discharge records at more than 400 US hospitals.

Any patient in the database who was discharged from a hospital in 2005 following a total abdominal hysterectomy was included in the study.

The subjects were classified into one of four treatment groups based on the clinical method used to close the surgical incision:

- Sutures
- Staples
- TSA
- Staples and TSA

While the study was open to all commercially available TSAs, at the time of the study the only TSA used on patients in the database was DERMABOND® Topical Skin Adhesive. Thus, the TSA group only had patients treated with DERMABOND Adhesive.

All treatment groups were assessed on three continuous outcomes: length of inpatient stay, total inpatient cost, and days of antibiotic treatment. Length of stay and inpatient cost was available directly from the database; antibiotic treatment days were estimated using the last date when a dose of antibiotic was administered.

Results

In total, 46,011 patients were included in the study. The method of wound closure for these patients is summarized in **Figure 1**.

Due to the large sample size, there were no statistically significant differences in the clinical, demographic, or hospital characteristics of the four treatment groups.

Figure 1
Distribution of Skin Closure Method

Skin Closure Method Evaluated in Study	# of Patients (n)
Sutures	21,201
Staples	23,441
TSA	880
Staples and TSA	489
All Methods	46,011

Length of Stay (LOS) and Total Costs

A summary of mean LOS and total hospitalization costs is shown in **Figure 2**.

While the difference in total costs between suture and TSA groups did not meet the significance requirement for this study ($P \leq 0.01$), the difference suggests lower total costs for the TSA group ($P = 0.039$).

Figure 2
LOS and Total Costs by Closure Method

Skin Closure Method Evaluated in Study	Mean LOS (days)	Mean Total Hospitalization Costs
Sutures	3.9	\$5,862
Staples	4.5	\$6,965
TSA	3.7	\$5,816
Staples and TSA	5.2	\$9,434

Conclusions

The results of this study demonstrate that the clinical and economic outcomes were consistently worse when staples were used to close an incision compared with use of suture or TSA alone.

The clinical outcomes resulting from the use of DERMABOND® Topical Skin Adhesive to close wounds were at least as good as the outcomes resulting from the use of suture to close wounds.

Additionally, there is evidence that the total costs of hospitalization for total hysterectomy patients may be less when the incision is closed with DERMABOND Adhesive versus sutures or staples.*

*This study was funded in full or in part by an educational grant from Ethicon, Inc.

Clinical Reference Article Summary

A Randomized Trial Comparing Octylcyanoacrylate Tissue Adhesive and Sutures In the Management of Lacerations

Quinn J, Wells G, Sutcliffe T, Jarmuske M, Maw J, Stiell I, Johns P

Source:

JAMA. 1997;277(19):1527-1530

Study Objective

The purpose of this study was to assess whether using DERMABOND® Topical Skin Adhesive for laceration repair is an effective alternative to suturing.

Methods

Patients with non-mucosal facial lacerations as well as certain extremity and torso lacerations, but not on hands, feet or joints, were eligible for this study.

Using a computer algorithm, patients were prospectively segregated into facial and non-facial groups and randomized into two groups—DERMABOND Adhesive and sutures.

In the suture group, lacerations were anesthetized and cleaned, as needed, before repair with a 5-0 or 6-0 monofilament suture. A dressing was applied for at least 48 hours.

In the DERMABOND Adhesive group, lacerations were cleaned with chlorhexidine and hemostasis was achieved using pressure or topical 1:1000 epinephrine. The wound edges were manually approximated and the adhesive was applied to the surface of the skin, covering the wound edges. The wound was held in place for 30 seconds. No dressing was applied.

The primary outcome was the cosmetic appearance of the scar, evaluated by a blinded plastic surgeon using a photograph of the wound taken 3 months after closure.

On two occasions, the surgeon examined the photograph and provided a cosmesis score based on a validated 100-mm visual analog scale, ranging from “best scar” to “worst scar.”

Additionally, time of procedure, patient pain, and wound complications (i.e., dehiscence, infection) were recorded. Time of procedure was evaluated from start of wound care to complete closure; patient pain and wound complications were recorded on a previously validated scale.

Wound complication was initially evaluated at 3-5 days for facial and at 10-14 days for torso and extremity lacerations. A second assessment occurred 3 months after closure.

Results

In total, 130 patients with 136 lacerations were included in the study. As summarized in **Figure 1**, an equal number of lacerations (68 per group) were randomized to the suture and DERMABOND Adhesive groups.

Figure 1
Patient Retention During Study

	DERMABOND Adhesive	Suture
Randomized	68	68
Initial follow-up	53	53
3 month follow-up	55	50
Withdrawn	1	1
Lost to follow-up	12	17
No Photographs	5	2
Completed Study	50	48

As shown in **Figure 2**, there was no significant difference in the blinded, 3-month cosmetic score of the DERMABOND® Topical Skin Adhesive group compared with the suture group. Similarly, there was no significant difference in wound complications between the suture group and the DERMABOND Adhesive group. Statistically significant differences were seen for patient pain and procedure time.

Figure 2
Summary of Observed Clinical Outcomes

	DERMABOND Adhesive	Suture	(P) Value
Mean Cosmetic Score (mm)	67	68	0.65
% Optimal Wound Scores (initial eval)	80%	82%	0.80
% Optimal Wound Scores (3 month eval)	72%	75%	0.74
Mean Pain Scores (mm)	7.2	18.0	<0.001
Mean Time of Procedure (min)	3.6	12.4	<0.001

Conclusions

The results of this study demonstrate that DERMABOND Adhesive produces cosmetic results similar to suturing on certain types of lacerations.

Additionally, lacerations closed with DERMABOND Adhesive were associated with shorter procedure time and less patient pain than lacerations closed with sutures.*

*This study was funded in full or in part by an educational grant from Ethicon, Inc.

For More Information

Call 1-877-ETHICON (384-4266)

In addition to support from Ethicon Sales Representatives, Ethicon's Medical Affairs team is available to provide balanced, non-promotional scientific information to healthcare professionals.

Medical information request form

To: Ethicon Medical Affairs

E-mail: Eth_Medical_Info@its.jnj.com Voicemail: (800) 888-9234, x3800

Date: _____

From (Requestor): _____

Name: _____

(Circle one):

M.D.

D.O.

R.N.

N.P.

Pharm.D.

Ph.D.

R.Ph.

Other: _____

Title: _____ Institution/Office: _____

Address: _____

City: _____ State: _____ ZIP: _____

Telephone: _____ Fax: _____

E-mail Address: _____

Desired Response Method (Circle one):

US Mail

Phone

E-mail

Fax

Meeting with Medical Affairs Representative

Requestor's Signature:

(REQUIRED FOR PROCESSING) _____

Please send medical information on the following topic(s):

(Be as specific as possible with respect to product topic, area of use, outcome of interest, etc.)

Sales Representative: _____ Territory: _____

PRINT FULL NAME _____

Bibliography

Listed below are all of the currently published RCTs that have evaluated the use of DERMABOND® Topical Skin Adhesive in an application consistent with the indication in the product's label (i.e., skin closure). Studies that evaluated the use of DERMABOND Adhesive for purposes inconsistent with the intended indication were excluded from the bibliography.

- Amin M, Glynn F, Timon C. Randomized trial of tissue adhesive vs staples in thyroidectomy integrating patient satisfaction and Manchester score. *Otolaryngol Head Neck Surg.* 2009;140(5):703-708.
- Blondeel, PNV, Murphy JW, Debrosse D, Nix III JC, Puls LE, Theodore N, Coulthard P. Closure of long surgical incisions with a new formulation of 2-octylcyanoacrylate tissue adhesive versus commercially available methods. *Am J Surg.* 2004;188(3):307-313.
- Brown JK, Campbell BT, Drongowski RA, Alderman AK, Geiger JD, Teitelbaum DH, Quinn J, Coran AG, Hirschl RB. A prospective, randomized comparison of skin adhesive and subcuticular suture for closure of pediatric hernia incisions: cost and cosmetic considerations. *J Pediatr Surg.* 2009;44(7):1418-1422.
- Bruns TB, Robinson BS, Smith RJ, Kile DL, Davis TP, Sullivan KM, Quinn JV. A new tissue adhesive for laceration repair in children. *J Pediatr.* 1998;132(6):1067-1070.
- Carleo C, Singer AJ, Thode HC Jr. Effect of frequent water immersion on the rate of tissue adhesive sloughing: a randomized study. *CJEM.* 2005;7(6):391-395.
- Chen K, Klapper AS, Voige H, Del Priore G. A randomized, controlled study comparing two standardized closure methods of laparoscopic port sites. *JSL.S.* 2010;14(3):391-394.
- Daykan Y, Sharon-Weiner M, Pasternak Y, et al. Skin closure at cesarean delivery, glue vs subcuticular sutures: A randomized controlled trial. *Am J Obstet Gyn.* 2017;216(4):406.e1-406.e5.
- El-Gazzar Y, Smith DC., Kim S.J., Hirsh DM, Blum Y, Cobelli M, Cohen HW. The Use of Dermabond® as an Adjunct to Wound Closure After Total Knee Arthroplasty: Examining Immediate Post-Operative Wound Drainage. *J Arthroplasty.* 2013;28:553-556.
- Eymann R, Kiefer M. Glue instead of stitches: a minor change of the operative technique with a serious impact on the shunt infection rate. *Acta Neurochir Suppl.* 2010;106:87-89.
- Gennari R, Rotmensz N, Ballardini B, Scevola S, Perego E, Zanini V, Costa A. A prospective, randomized, controlled clinical trial of tissue adhesive (2-octylcyanoacrylate) versus standard wound closure in breast surgery. *Surgery.* 2004;136(3):593-599.
- Glennie RA, Korczak A, Naudie DD, et al. MONOCRYL and DERMABOND vs staples in total hip arthroplasty performed through a lateral skin incision: A randomized controlled trial using a patient-centered assessment tool. *J Arthroplasty.* 2017;32(8):2431-2435.
- Greene D, Koch RJ, Goode RL. Efficacy of octyl-2-cyanoacrylate tissue glue in blepharoplasty. A prospective controlled study of wound-healing characteristics. *Arch Facial Plast Surg.* 1999;1(4):292-296.
- Handschel JG, Depprich RA, Dirksen D, Runte C, Zimmermann A, Kübler NR. A prospective comparison of octyl-2- cyanoacrylate and suture in standardized facial wounds. *Int J Oral Maxillofac Surg.* 2006;35(4):318-323.
- Holger JS, Wandersee SC, Hale DB. Cosmetic outcomes of facial lacerations repaired with tissue-adhesive, absorbable, and nonabsorbable sutures. *Am J Emerg Med.* 2004;22(4):254-257.
- Hollander JE, Singer AJ. Application of tissue adhesives: rapid attainment of proficiency. *Acad Emerg Med.* 1998;5(10):1012-1017
- Jallali N, Haji A, Watson CJ. A prospective randomized trial comparing 2-octyl cyanoacrylate to conventional suturing in closure of laparoscopic cholecystectomy incisions. *J Laparoendosc Adv Surg Tech A.* 2004;14(4):209-211.
- Kent A, Liversedge N, Dobbins B, McWhinnie D, Jan H. A prospective, randomized, controlled, double-masked, multi-center clinical trial of medical adhesives for the closure of laparoscopic incisions. *J Minim Invasive Gynecol.* 2014;21:252-258.
- Khan RJ, Fick D, Yao F, Tang K, Hurworth M, Nivbrant B, Wood D. A comparison of three methods of wound closure following arthroplasty: a prospective, randomised, controlled trial. *J Bone Joint Surg Br.* 2006;88(2):238-242.
- Koonce SL, Eck DL, Shaddix KK, Perdakis G. A prospective randomized controlled trial comparing N-butyl-2 cyanoacrylate (Histoacryl), octyl cyanoacrylate (Dermabond), and subcuticular suture for closure of surgical incisions. *Ann Plast Surg.* 2015;74:107-110.
- Krishnamoorthy B, Najam O, Khan UA, Waterworth P, Fildes JE, Yonan N. Randomized prospective study comparing conventional subcuticular skin closure with Dermabond skin glue after saphenous vein harvesting. *Ann Thorac Surg.* 2009;88(5):1445-1449.
- Maartense S, Bemelman WA, Dunker MS, de Lint C, Pierik EG, Busch OR, Gourma DJ. Randomized study of the effectiveness of closing laparoscopic trocar wounds with octylcyanoacrylate, adhesive papertape or polyglactone. *Br J Surg.* 2002;89(11):1370-1375.
- Maloney J, Rogers GS, Kapadia M. A prospective randomized evaluation of cyanoacrylate glue devices in the closure of surgical wounds. *J Drugs Dermatol.* 2013;12:810-814.
- Man SY, Wong EM, Ng YC, Lau PF, Chan MS, Lopez V, Mak PS, Graham CA, Rainer TH. Cost-consequence analysis comparing 2-octyl cyanoacrylate tissue adhesive and suture for closure of simple lacerations: A randomized controlled trial. *Ann Emerg Med.* 2009;53(2):189-197.
- Martin JG, Hollenbeck ST, Janas G, Makar RA, Pabon-Ramos WM, Suhocki PV, Miller MJ, Sopko DR, Smith TP, Kim CY. Randomized controlled trial of octyl cyanoacrylate skin adhesive versus subcuticular suture for skin closure after implantable venous port placement. *J Vasc Interv Radiol.* 2017;28:111-116.
- Matin SF. Prospective randomized trial of skin adhesive versus sutures for closure of 217 laparoscopic port-site incisions. *J Am Coll Surg.* 2003 Jun;196(6):845-853.
- Mattick A, Beattie T, Ahmad T. A randomised, controlled trial comparing a tissue-sensitive (2-octylcyanoacrylate) with adhesive strips (Steristrips) for paediatric laceration repair. *Emerg Med J.* 2002;19(5): 405-407.
- Mota R, Costa F, Amaral A, Oliveira F, Santos CC, Ayres-De-Campos D. Skin adhesive versus subcuticular suture for perineal skin repair after episiotomy - A randomized controlled trial. *Acta Obstet Gynecol Scand.* 2009;88(6):660-666.
- Mudd CD, Boudreau JA, Moed BR. A prospective randomized comparison of two skin closure techniques in acetabular fracture surgery. *J Orthop Traumatol.* 2014;15:189-194.

Nipshagen MD, Hage JJ, Beekman WH. Use of 2-octyl-cyanoacrylate skin adhesive (Dermabond) for wound closure following reduction mammoplasty: A prospective, randomized intervention study. *Plast Reconstr Surg*. 2008;122(1):10-18.

Ong CC, Jacobsen AS, Joseph VT. Comparing wound closure using tissue glue versus subcuticular suture for pediatric surgical incisions: A prospective, randomised trial. *Pediatr Surg Int*. 2002;18(5-6):553-555.

Ong J, Ho KS, Chew MH, Eu KW. Prospective randomised study to evaluate the use of DERMABOND ProPen (2- octylcyanoacrylate) in the closure of abdominal wounds versus closure with skin staples in patients undergoing elective colectomy. *Int J Colorectal Dis*. 2010;25(7):899-905.

Osmond MH, Quinn JV, Sutcliffe T, Jarmuske M, Klassen TP. A randomized, clinical trial comparing butylcyanoacrylate with octylcyanoacrylate in the management of selected pediatric facial lacerations. *Acad Emerg Med*. 1999;6(3):171-177.

Patel HM, Shah MJ, Duttaroy DD, Kacheriwala SM, Patel SJ, Patel RM. Superiority of octyl-2 cyanoacrylate over polyamide black for surgical site incisions. Prospective randomized trial. *Surg Chron*. 2013; 18(3):139-143.

Pronio A, Di Filippo A, Narilli P, Caporilli D, Vestri A, Ciamberlano B, Pelle F, Montesani C. Closure of cutaneous incision after thyroid surgery: A comparison between metal clips and cutaneous octyl-2-cyanoacrylate adhesive. A prospective randomized clinical trial. *Eur J Plast Surg*. 2010;34(2):103-110.

Quinn J, Wells G, Sutcliffe T, Jarmuske M, Maw J, Stiell I, Johns P. A randomized trial comparing octylcyanoacrylate tissue adhesive and sutures in the management of lacerations. *JAMA*. 1997;277(19):1527-1530.

Ridgway DM, Mahmood F, Moore L, Bramley D, Moore PJ. A blinded, randomised, controlled trial of stapled versus tissue glue closure of neck surgery incisions. *Ann R Coll Surg Engl*. 2007;89(3):242-246.

Romero P, Frongia G, Wingerter S, Holland-Cunz S. Prospective, randomized, controlled trial comparing a tissue adhesive (Dermabond) with adhesive strips (Steri-Strips) for the closure of laparoscopic trocar wounds in children. *Eur J Pediatr Surg*. 2011;21(3):159-162.

Sebesta MJ, Bishoff JT. Octylcyanoacrylate skin closure in laparoscopy. *JSL*. 2004;8(1):9-14.

Shamiyeh A, Schrenk P, Stelzer T, Wayand WJ. Prospective randomized blind controlled trial comparing sutures, tape, and octylcyanoacrylate tissue adhesive for skin closure after phlebectomy. *Dermatol Surg*. 2001;27(10):877-880.

Siddiqui M, Bidaye A, Baird E, Abu-Rajab R, Stark A, Jones B, Ingram R, Anthony I. Wound dressing following primary total hip arthroplasty: A prospective randomised controlled trial. *J Wound Care*. 2016;25-40.

Singer AJ, Giordano P, Fitch JL, Gulla J, Ryker D, Chale S. Evaluation of a new high-viscosity octylcyanoacrylate tissue adhesive for laceration repair: a randomized, clinical trial. *Acad Emerg Med*. 2003;10(10):1134-1137.

Singer AJ, Hollander JE, Valentine SM, Turque TW, McCuskey CF, Quinn JV. Prospective, randomized, controlled trial of tissue adhesive (2- octylcyanoacrylate) vs standard wound closure techniques for laceration repair. *Acad Emerg Med*. 1998;5(2):94-99.

Singer AJ, Quinn JV, Clark RE, Hollander JE. Closure of lacerations and incisions with octylcyanoacrylate: A multicenter randomized controlled trial. *Surgery*. 2002;131(3):270-276.

Sniezek PJ, Walling HW, DeBloom JR 3rd, Messingham MJ, VanBeek MJ, Kreiter CD, Whitaker DC, Arpey CJ. A randomized controlled trial of high-viscosity 2-octyl cyanoacrylate tissue adhesive versus sutures in repairing facial wounds following Mohs micrographic surgery. *Dermatol Surg*. 2007;33(8):966-971.

Soni A, Narula R, Kumar A, Parmar M, Sahore M, Chandel M. Comparing cyanoacrylate tissue adhesive and conventional subcuticular skin sutures for maxillofacial incisions - A prospective randomized trial considering closure time, wound morbidity, and cosmetic outcome. *J Oral Maxillofac Surg*. 2013 Dec;71(12):2152.e1-8.

Spencer S, Coban N, Koch L, Schirdewan A, Mueller D. Comparison of skin adhesive and absorbable intracutaneous suture for the implantation of cardiac rhythm devices. *Europace*. 2011;13:416-420.

Strauss EJ, Weil WM, Jordan C, Paksima N. A prospective, randomized, controlled trial of 2-octylcyanoacrylate versus suture repair for nail bed injuries. *J Hand Surg Am*. 2008;33(2):250-253.

Sun J, Chen Q-M, Zhang M, Shi C-R. Octylcyanoacrylate versus absorbable suture in the repair of skin wound in children. *Chinese J Clin Rehabilitation*. 2005;19:26-29.

Switzer EF, Dinsmore RC, North JH Jr. Subcuticular closure versus dermabond: A prospective randomized trial. *Am Surg*. 2003;69(5):434-436.

Tierney EP, Moy RL, Kouba DJ. Rapid absorbing gut suture versus 2 octylethylcyanoacrylate tissue adhesive in the epidermal closure of linear repairs. *J Drugs Dermatol*. 2009;8(2):115-119.

Toriumi DM, O'Grady K, Desai D, Bagal A. Use of octyl-2-cyanoacrylate for skin closure in facial plastic surgery. *Plast Reconstr Surg*. 1998;102(6):2209-2219.

Wong EM, Rainer TH, Ng YC, Chan MS, Lopez V. Cost-effectiveness of Dermabond versus sutures for lacerated wound closure: A randomised controlled trial. *Hong Kong Med J*. 2011;17:4-8.

Zempsky WT, Parotti D, Grem C, Nichols J. Randomized controlled comparison of cosmetic outcomes of simple facial lacerations closed with Steristrip closures or Dermabond tissue adhesive. *Pediatr Emerg Care*. 2004;20(8):519-524.